



DID YOU KNOW?

RSV is a leading cause of hospitalization in the US for children <1 year of age



In these uncertain times,

You have the Power to help Protect

Protect your vulnerable baby from severe Respiratory Syncytial Virus (RSV) disease

IMPORTANT



SCHEDULE MONTHLY SYNAGIS DOSING APPOINTMENTS every 28-30 days with your baby's doctor throughout RSV season.

RSV season typically occurs from late fall through spring.*

IMPORTANT SAFETY INFORMATION

Who should not receive SYNAGIS?

Children should not receive SYNAGIS if they have ever had a severe allergic reaction to it. Signs and symptoms of a severe allergic reaction could include itchy rash; swelling of the face; difficulty swallowing; difficulty breathing; bluish color of the skin; muscle weakness or floppiness; and/or unresponsiveness. If your child has any of these signs or symptoms of a severe allergic reaction after getting SYNAGIS, call your child's healthcare provider or get medical help right away.

All imagery is for illustrative purposes only.

*RSV season typically lasts 5 months in the United States, but can vary by year and geographic region. High-risk infants should receive monthly doses (every 28-30 days) throughout RSV season.

[Please click here for additional Important Safety Information and full Prescribing Information, including Patient Information.](#)






Help protect your baby with SYNAGIS


Financial help may be available

Commercially eligible patients may get up to **\$6000** per SYNAGIS season to assist with out-of-pocket costs for SYNAGIS (paying as little as **\$0** per dose)

To join the SYNAGIS Copay Program[†]:

Talk to your doctor's office or Specialty Pharmacy  Your doctor or Specialty Pharmacy can enroll you in this program

OR

Enroll in the SYNAGIS Copay Program  Visit **SYNAGIS CONNECT™**

SYNAGIS CONNECT™ is a patient support program created by Sobi to provide individualized support to help appropriate patients get access to SYNAGIS.

You can get assistance with out-of-pocket costs, understanding your insurance coverage for SYNAGIS, and support resources. If you have any questions or for additional assistance, call **SYNAGIS CONNECT™** at 1-833-SYNAGIS (1-833-796-2447), Monday through Friday from 8:00 AM to 8:00 PM EST.

You will also find more information about protecting your baby with SYNAGIS on SYNAGIS.com

[†]Eligibility requirements apply. Patient must be a resident of the US or Puerto Rico. Patient must be commercially insured. There are no income requirements to participate in the program. Patient must not be insured by any government, state, or federally funded prescription program, including Medicare, Medicaid, Medigap, VA, DOD, or TRICARE.

IMPORTANT SAFETY INFORMATION (continued)

How is SYNAGIS given?

SYNAGIS is given as a monthly injection, usually in the thigh (leg) muscle, by your child's healthcare provider. If your child has a problem with bleeding or bruises easily, an injection could cause a problem. Your child should receive their first injection of SYNAGIS before the RSV season starts, to help protect them before RSV becomes active. RSV season is usually fall through spring, but it may begin earlier or last longer in certain areas. When RSV is most active, your child will need to receive injections of SYNAGIS every 28-30 days to help protect them from severe RSV disease for about a month. Your child should continue to receive monthly injections of SYNAGIS until the end of RSV season. Your child may still get severe RSV disease after receiving SYNAGIS. If your child has an RSV infection, they should continue to get their monthly injections throughout the RSV season to help prevent severe disease from new RSV infections.

The effectiveness of injections of SYNAGIS given less than monthly throughout the RSV season has not been established.

What are the possible side effects of SYNAGIS?

Serious side effects include severe allergic reactions, which may happen after any injection of SYNAGIS and may be life-threatening or cause death. Call your child's healthcare provider or get medical help right away if your child has any of the signs or symptoms of a serious allergic reaction. See "Who should not receive SYNAGIS?" for more information.

Common side effects of SYNAGIS include fever and rash.

These are not all the possible side effects of SYNAGIS.

APPROVED USE

SYNAGIS is a prescription medication that is used to help prevent a serious lung disease caused by respiratory syncytial virus (RSV) in children:

- born prematurely (at or before 35 weeks) **and** who are 6 months of age or less at the beginning of RSV season
- who have a chronic lung condition called bronchopulmonary dysplasia (BPD), that needed medical treatment within the last 6 months, **and** who are 24 months of age or less at the beginning of RSV season
- born with certain types of heart disease **and** who are 24 months of age or less at the beginning of RSV season

It is not known if SYNAGIS is safe and effective:

- to *treat* the symptoms of RSV in a child who already has RSV. SYNAGIS is used to help *prevent* RSV disease
- in children who are older than 24 months of age at the start of dosing

[Please click here for full Prescribing Information, including Patient Information.](#)

 Learn more about us at SOBI.com



SYNAGIS® is a registered trademark of Arexis AB c/o Swedish Orphan Biovitrum AB (publ)
SYNAGIS CONNECT™ is a trademark of Arexis AB c/o Swedish Orphan Biovitrum AB (publ)
© 2021 Swedish Orphan Biovitrum. All rights reserved. PP-10357 02/21

